

Helicobacter pylori, antigen (F - HepyAg)

Certain medicinal substances may affect the test result (see the table below). Antibiotic treatment and helicobacter eradication treatment should be discontinued at least four weeks before the test. The use of proton pump inhibitors (PPIs) should be discontinued at least 2 weeks before the test.

- Collect some stool into a clean bowl.
- Use the spoon on the lid of the sample container to collect the sample (tablespoonful, no more than half the volume of the sample container). Close the container carefully.
- Write down your name, personal identity number and the date of collection on a label sticker. Attach the sticker onto the side of the container. Place the container into a sample bag and close the bag. Carefully fill out all the fields in this form.
- Deliver the sample and this completed form to the laboratory where you received the sampling supplies.

Please fill in carefully the following fields:

Name: _____

Social security number/date of birth: _____

Date of collection: ____/____/____

MEDICATION	DISCONTINUATION TIME BEFORE TESTING
Helicobacter eradication treatment	Minimum four weeks
Antibiotics	Minimum four weeks
<u>Proton pump inhibitors (PPI):</u>	Minimum two weeks
Esomeprazole	Acidal, Esomeprazol Actavis, Esomeprazol Krka, Esomeprazol Orion, Esomeprazole Orifarm, Esomeprazol Ratiopharm, eEsomeprazole Sandoz, Nexium,
Lansoprazole	Vimovo
Omeprazole	Gasterix, Helipak, Lansoprazol, Zolt
Pantoprazole	Losec, Losec Mups, Omeprazol Actavis, Omeprazol Sanoswiss, Omeprazol Ratiopharm, Omestad
Rabeprazole	Pantoprazol Actavis, Pantoprazol Krka, Pantoprazol Mylan, Pantoprazol Orion, Pantoprazole Ratiopharm, Pantoprazol Sandoz, Somac control, Somac, Pariet, Rabeprazol Krka
Bismuth:	Minimum two weeks
	Gastrodenol, Pylera

Inquiries: